

$$a_1, a_2, a_3, \dots, a_n$$

$$a_1 = 2.5$$

$$: 17 - \quad 80 - \quad 33 -$$

$$a_{33} = a_{17} + 80$$

$$a_1 + 32d = a_1 + 16d + 80$$

$$16d = 80$$

$$d = 5$$

:

$$a_3, a_6, a_9, \dots, a_n$$

$$3 -$$

$$, (\quad)$$

$$, k - \quad , a_{k+3} - a_k = a_k + 3d - a_k = 3d = 3 \cdot 5 = 15$$

$$a_6 - a_3 = a_3 + 3d - a_3 = 3d = 3 \cdot 5 = 15$$

$$. 15$$

$$. 3100$$

$$a_3 = a_1 + 2d = 2.5 + 2 \cdot 5 = 12.5 - , t = \frac{n}{3} , \quad t - \quad (1)$$

$$S_n = \frac{n(2a_1 + d(n-1))}{2}$$

$$3100 = \frac{t(2 \cdot 12.5 + 15(t-1))}{2}$$

$$6200 = t(25 + 15t - 15)$$

$$6200 = t(15t + 10)$$

$$0 = 15t^2 + 10t - 6200$$

$$t_{1,2} = \frac{-10 \pm 610}{30}$$

$$t = 20 \quad \leftarrow t > 0$$


$$. 20 \quad :$$

$$. 60 \quad , \quad (2)$$

$$. 60 \quad :$$

, AC = AB ,

, AD -



$$CD = BD = \frac{26}{2} = 13 \text{ cm}$$

$$\sphericalangle DAC = \sphericalangle DAB = \frac{64^\circ}{2} = 32^\circ$$

ACD

$$\tan 32^\circ = \frac{CD}{AD}$$

$$AD = \frac{13}{\tan 32^\circ}$$

$$\boxed{AD = 20.80 \text{ cm}}$$

$$.S = \frac{BC \cdot AD}{2} = \frac{26 \cdot 20.80}{2} = 270.46 \text{ cm}^2 :$$

. " 8112

$$8112 = 270.46 \cdot H \rightarrow H = 29.99 \approx 30 \text{ cm}$$

. " 30

, ABC

() A'B

$\sphericalangle A'BA$

, ABC

A'B

. ($\sphericalangle A'AB = 90^\circ$) A'AB

A'AB

$$\tan \sphericalangle A'BA = \frac{AA'}{AB}$$

$$\tan \sphericalangle A'BA = \frac{30}{24.53}$$

$$\boxed{\sphericalangle A'CA = 50.73^\circ}$$

. 50.73° ABC

ABD

$$\sin 32^\circ = \frac{BD}{AB}$$

$$AB = \frac{13}{\sin 32^\circ}$$

$$\boxed{AB = 24.53 \text{ cm}}$$

A'B

. ($\sphericalangle AA'D' = 90^\circ$) AA'D'

A'AD'

AA'D'

$$\tan \sphericalangle A'AD' = \frac{A'D'}{AA'}$$

$$\tan \sphericalangle A'AD' = \frac{20.80}{30}$$

$$\boxed{\sphericalangle A'AD' = 34.73^\circ}$$

. 34.73° A'AD'

"

$$y = 0.5x \quad f(x) = 0.5x - 0.5 \cos(2x) + 0.5$$

$$-f \leq x \leq f \quad x \quad (1)$$

$$0.5x = 0.5x - 0.5 \cos(2x) + 0.5$$

$$0.5 \cos 2x = 0.5$$

$$\cos 2x = 1$$

$$2x = 2fk$$

$$x = fk$$

$$x = 0, f, -f \quad k = 0, 1, -1$$

$$x = 0, f, -f :$$

0.5

(2)

$$f'(x) = 0.5 + \sin 2x$$


$$f'(0) = 0.5 + \sin(2 \cdot 0) = 0.5 \quad o.k.$$

$$f'(f) = 0.5 + \sin(2 \cdot f) = 0.5 \quad o.k.$$

$$f'(-f) = 0.5 + \sin(2 \cdot (-f)) = 0.5 \quad o.k.$$

:

$$-f \leq x \leq f$$



()

$$S = \int_{-f}^f (0.5x - 0.5 \cos(2x) + 0.5 - 0.5x) dx$$

$$S = \int_{-f}^f (-0.5 \cos(2x) + 0.5) dx$$

$$S = \left(-\frac{\sin 2x}{4} + 0.5x \right) \Big|_{-f}^f$$

$$S = \left(-\frac{\sin 2f}{4} + 0.5f \right) - \left(-\frac{\sin(2(-f))}{4} + 0.5(-f) \right)$$

$$S = 0.5f + 0.5f$$

$$S = f$$

f :

(0 - m) $f(x) = 2x^2 e^{-\frac{x^2}{m}}$.
 . x :
 , x = -2 - .

$$f'(x) = 4xe^{-\frac{x^2}{m}} + 2x^2 e^{-\frac{x^2}{m}} \left(-\frac{2x}{m}\right) = 2e^{-\frac{x^2}{m}} (2x - x^2 \cdot \frac{2x}{m})$$

$$0 = 2 \cdot (-2) - (-2)^2 \cdot \frac{2 \cdot (-2)}{m}$$

$$0 = -4 + \frac{16}{m} \rightarrow 0 = -4m + 16 \rightarrow \boxed{m = 4}$$

. m = 4 :

((- f(-x) = f(x)) f(x) = 2x^2 e^{-\frac{x^2}{4}} m = 4)
 (0, 0) 2x^2 = 0 y = 0 x - (1) .
 . y -

. (0, 0) :

: (2)

$$f'(x) = 4xe^{-\frac{x^2}{4}} + 2x^2 e^{-\frac{x^2}{4}} \left(-\frac{2x}{4}\right)$$

$$\boxed{f'(x) = 2e^{-\frac{x^2}{4}} (2x - 0.5x^3)}$$

$$0 = x(2 - 0.5x^2) \rightarrow x = 0 \rightarrow (0, 0)$$

$$2 - 0.5x^2 = 0 \rightarrow x^2 = 4 \rightarrow x = \pm 2$$

$$x = 2 \rightarrow f(2) = 2 \cdot 2^2 e^{-\frac{2^2}{4}} \rightarrow \boxed{\left(2, \frac{8}{e}\right)} \rightarrow \boxed{\left(-2, \frac{8}{e}\right)}$$


$$\left(2e^{-\frac{x^2}{m}} \right)$$

$$f'(-3) = 2 \cdot (-3) - 0.5(-3)^3 > 0, \quad f'(-1) < 0, \quad f'(1) > 0, \quad f'(3) < 0$$

	-2		0		2		x
+	0	-	0		0	-	y'
↖	Max	↘	Min	↖	Max	↘	


. (0, 0) , (-2, $\frac{8}{e}$) , (2, $\frac{8}{e}$) :

(3)



$x < 0$ $0 < x < 2$ $f'(x) > 0$ - , $f(x)$, / .
- $-2 < x < 0$ $x > 2$ $f'(x) < 0$ --
- $f'(x) = 0$, $-2 \leq x \leq 2$

:



$$f(x) = -x \ln(2x)$$

$$OC \cdot OA$$

$$f(x) = 0, x =$$

C

$$0 = -x \ln(2x)$$

$$x = 0 \rightarrow x_0 = 0$$

$$\ln(2x) = 0 \rightarrow 2x = e^0 = 1 \rightarrow x_c = 0.5$$

$$OC = 0.5 - 0 = 0.5 :$$

AB

$$0 < x < 0.5$$

$$f'(x) = -\ln(2x) - x \cdot \frac{2}{2x}$$

$$\boxed{f'(x) = -\ln(2x) - 1}$$

$$0 = -\ln(2x) - 1$$

$$\ln(2x) = -1$$

$$2x = e^{-1} = \frac{1}{e}$$

$$x = \frac{1}{2e} \rightarrow f\left(\frac{1}{2e}\right) = -\frac{1}{2e} \cdot \ln\left(2 \cdot \frac{1}{2e}\right) = \frac{1}{2e}$$

$$OA = \frac{1}{2e} - 0 = \frac{1}{2e}, y = \frac{1}{2e} \quad AB$$

$$OC \cdot OA = 0.5 \cdot \frac{1}{2e} = \frac{1}{4e} :$$

$$\frac{1}{4e} :$$

35805

13

$$- M_0 \quad , \quad M_t = M_0 \cdot q^t :$$

$$.t \quad M_t , \quad q$$

80% - , 20% - 10

$$. \quad 10 - 0.6M_0 - M_0 -$$

$$0.8M_0 = M_0 \cdot q^{10} \quad / : M_0$$

$$0.8 = q^{10}$$

$$q = \sqrt[10]{0.8}$$

$$\boxed{q = 0.9779}$$

40% - ,

60% -

$$0.6M_0 = M_0 \cdot 0.9779^t \quad / : M_0$$

$$0.6 = 0.9779^t$$

$$\ln 0.6 = \ln 0.9779^t$$

$$\ln 0.6 = t \ln 0.9779$$

$$\frac{\ln 0.6}{\ln 0.9779} = t$$

$$\boxed{t = 22.89}$$

22.89